

(18)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 068 590
B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 02.01.85

(61) Int. Cl.: **C 07 C 127/17,**
C 07 C 157/07, A 61 K 31/17

(21) Application number: 82200796.9

(22) Date of filing: 28.06.82

(54) Novel urea derivatives.

(30) Priority: 30.06.81 US 278970
25.05.82 US 381672

(43) Date of publication of application:
05.01.83 Bulletin 83/01

(45) Publication of the grant of the patent:
02.01.85 Bulletin 85/01

(84) Designated Contracting States:
BE DE FR GB IT NL

(68) References cited:
DE-A-2 942 564
GB-A-1 153 261
JP-A-49 013 785
US-A-2 663 730

(70) Proprietor: **THE PROCTER & GAMBLE**
COMPANY
301 East Sixth Street
Cincinnati Ohio 45202 (US)

(72) Inventor: **Buckwalter, Brian Lee**
621 Glensprings Dr.
Springdale Ohio 45246 (US)
Inventor: **Lahann, Thomas Robert**
9239 Harrison Pike
Cleveland Ohio 45002 (US)

(74) Representative: **Suslic, Lydia et al**
Procter & Gamble European Technical Center
Temselaan 100
B-1820 Strombeek-Bever (BE)

EP 0 068 590 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).

Courier Press, Leamington Spa, England.

Description

Technical Field

This invention provides certain novel urea derivatives having analgesic and anti-irritant activity.

In general, analgesics fall into two broad categories. The simple analgesics, such as aspirin, are most effective against pain of integumental origin, headache, and muscle aches; the narcotics are more useful for deep or visceral pain. Narcotic analgesics such as morphine also produce more profound effects than simple analgesics, and are potentially addicting, with the development of tolerance and physical dependence. The narcotic analgesics appear to work through interaction with the endorphin/enkephalin system of the CNS; many of the simple, non-narcotic analgesics appear to work by inhibition of prostaglandin synthetase. The effect of narcotics is to elevate the pain threshold above the normal level; the non-narcotic analgesics act to raise an abnormally low pain threshold to the normal level. The narcotic analgesics are antagonized by N-allyl compounds such as naloxone; the non-narcotic analgesics are not.

In the field of anti-irritancy, dermal and tissue irritation can be caused by common irritants such as acids and alkalis, keratolytics and proteolytics, depilatories, plant oils, and the like. In many instances, it is observed that compounds which effectively block the action of one class of irritants are totally or substantially ineffective with other classes of irritants. The problem in the art has been compounded by the inadequacy, or unavailability in some cases, of appropriate animal models of dermal irritation. For example, most laboratory animals are insensitive to poison ivy. As a consequence, the search has long continued, in a purely empirical way, for anti-irritant compositions — not always successfully.

The present invention relates to the discovery that certain novel urea derivatives are analgesics and anti-irritants. In their analgesic action, these compounds appear to be largely unrelated to the two known classes of analgesics. In certain tests, these compounds produce a level of analgesia comparable to morphine, yet do not appear to involve the endorphin-enkephalin system, and thus should not be reversed by narcotic antagonists, such as naloxone. It is believed that these compounds will effectively prevent the development of cutaneous hyperalgesia. At high doses, it is believed that these compounds will also exert analgesic activity in classical models of deep pain, elevating the pain threshold above the normal value.

In their anti-irritant activity, these compounds are active against several irritants, including croton oil, and particularly depilatories (thioglycolates). They appear to act, not as protectants or barriers, but, rather, directly to block or eliminate responses to the irritant chemical. In addition, the urea derivatives of this invention offer the benefit of reduced discomfort on application. By contrast, compounds such as capsaicin, which possess potent anti-irritant activity, can cause irritation and reddening of the skin.

Background Art


J. A. Kleman, *Quart. J. of Exp. Physiol.*, (1972) 62, 151—161, states that capsaicin, N-(3-methoxy-4-hydroxybenzyl)-8-methyl-6-nonenamide, is known to confer resistance to certain chemical irritants.

Jancso, et al., *Br. J. Pharmac. Chemother.* (1967), 31, 138—151, states that by repeated administration of capsaicin, ocular and/or cutaneous pain receptors can be desensitized to chemical, but not other, stimuli.

Arvier, et al., *Br. J. Pharm.* (1977), 59, 61—68, indicate that capsaicin reduces or blocks the edema formation associated with certain types of inflammation.

Disclosure of the Invention

The present invention provides compounds of the following formula:



wherein X is O or S; wherein n = 1 or 0; wherein R is C₃—C₁₂ (preferably C₈—C₁₂) alkyl or aralkyl, branched or unbranched, aryl or haloaryl, or C₂—C₂₂ (preferably C₆—C₂₂) alkenyl, branched or unbranched; wherein R₁ is H, OH or OCH₃ and R₂ is H or OH (at least one of R₁ and R₂ being OH or OCH₃) when n = 1, and wherein R₁ is OCH₃ or OH and R₂ is OH when n = 0; and pharmaceutically acceptable salts thereof. These compounds or salts thereof are hereafter generally referred to as urea derivatives. The most preferred urea derivatives are those where X is S, where n = 1, where R₁ is OCH₃ and where R₂ is OH (the N-(3-methoxy-4-hydroxybenzyl) thioureas), and especially those where R is n-

octyl, phenyl or 1,1,3,3-tetramethylbutyl. These compounds can be made by synthesis from readily available starting materials, using standard synthetic techniques, as described hereinafter. In the practice of this invention, these urea derivatives can be administered either topically or systemically.

5 A. Definitions

By "safe and effective amount" is meant an amount of the urea derivative sufficient to provide analgesic or anti-irritant activity so as to alleviate or prevent the pain or irritation being treated at a reasonable benefit/risk ratio attendant with any medical treatment. The amount of the urea derivative used will vary with the particular condition being treated, the severity of the condition, the duration of
10 the treatment, the specific formulation employed and the concentration of the urea derivative therein, and like factors.

By "systemic administration" is meant introducing the urea derivative or composition containing same into the tissues of the body, other than by topical application. Systemic administration thus includes, without limitation, oral, intrathecal, epidural, intramuscular, intravenous, intraperitoneal, and
15 subcutaneous administration.

By "pharmaceutically acceptable" salts is meant those salts which are safe for topical or systemic administration. These include the sodium, potassium, calcium, magnesium, and ammonium salts.

By "topical application" herein is meant directly laying on or spreading the urea derivative, or composition containing same, on epidermal or epithelial tissue (including outer skin and oral, gingival,
20 nasal, etc. tissue).

By the term "comprise" as used herein is meant that various other inert ingredients, compatible drugs and medicaments, and steps can be employed in the compositions and processes of this invention as long as the critical urea derivatives are present in the compositions and are used in the manner disclosed. The term "comprising" thus encompasses and includes the more restrictive terms
25 "consisting essentially of" and "consisting of" which characterize the use of the essential urea derivatives in the compositions and methods disclosed herein.

By "compatible" herein is meant that the components of the composition are capable of being commingled without interacting in a manner which would substantially decrease the efficacy of the total compositions under ordinary use situations.

30 All percentages herein are by weight of the composition unless otherwise specified.

B. Synthesis of the Urea Derivatives of the Present Invention

The urea derivatives of this invention can readily be prepared by synthesis along the lines of the following schematic and illustrative examples of the N-(3-methoxy-4-hydroxybenzyl and phenyl) ureas
35 and thioureas.

40

45


50

55

60

65

0 068 590



Example I

N-(3-methoxy-4-hydroxybenzyl)-N'-octylurea was prepared by the following method.

1. 3-methoxy-4-(2-ethoxy-1-oxapropyl)-benzonitrile

2 ml of trifluoroacetic acid was added slowly to a chilled solution of 20 g of 3-methoxy-4-hydroxybenzonitrile, 50.7 ml of ethyl vinyl ether and 50 ml tetrahydrofuran (THF). The mixture was stirred at room temperature overnight. 7.5 ml of triethylamine was added to the mixture and stirring was continued for several minutes until the solvent was evaporated. The residue was partitioned between 400 ml of diethyl ether and 100 ml 1N NaOH. The diethyl ether layer was washed successively with 100 ml H₂O and 100 ml brine, dried over MgSO₄ and evaporated to yield 31.2 gm of 3-methoxy-4-(2-ethoxy-1-oxapropyl)-benzonitrile.

2. 2-methoxy-4-aminomethyl-(2-ethoxy-1-oxapropyl)benzene

A solution of 5.0 g 3-methoxy-4-(2-ethoxy-1-oxapropyl)benzonitrile in 25 ml anhydrous THF was added dropwise to 119.5 ml 1 M lithium aluminum hydride in THF. The mixture was heated at reflux for four hrs, cooled to 0°, and the excess reagent quenched with 5 N NaOH. Precipitated aluminum salts were removed by filtration and the THF was evaporated. The residue was partitioned between 200 ml ether and 75 ml H₂O. The ether layer was washed with 75 ml of brine, dried over K₂CO₃, filtered, and evaporated to yield 3.63 g of 2-methoxy-4-aminomethyl-(2-ethoxy-1-oxapropyl)benzene.


3. N-(3-methoxy-4-(2-ethoxy-1-oxapropylbenzyl))-N'-octylurea

A solution of 2.92 g of octyl isocyanate in 25 ml diethyl ether was added dropwise to a stirred solution of 3.63 g of 2-methoxy-4-aminomethyl-(2-ethoxy-1-oxapropyl)benzene in 25 ml diethyl ether cooled in an ice bath over a 30 minute period. The reaction mixture was allowed to stir four hours at room temperature. The solvent was evaporated to yield 6.66 g of the crude urea which was purified by column chromatography on silica gel with 60% ethyl acetate-hexane, yielding 5.2 g of the urea.

4. N-(3-methoxy-4-hydroxybenzyl)-N'-octylurea

30 ml 1 N HCl was added to an ice cold solution of the urea in 50 ml of THF and the mixture was stirred for 45 minutes. The THF was removed *in vacuo* and the residue was partitioned between 100 ml CH₂Cl₂ and 100 ml H₂O. The organic phase was then washed with 50 ml H₂O and 50 ml brine, dried over K₂CO₃, filtered and evaporated. The crude product was recrystallized from CH₂Cl₂ pentane which yielded 3.07 gm of white crystals, M.P. 90—92°C. Analysis calculated for C₁₇H₂₈N₂O₃: C, 66.20; H, 9.15; N, 9.08. Found: C, 66.27; H, 9.22; N, 8.98.

The thioureas can be prepared more directly, because the less reactive isothiocyanates selectively attack the amino group, obviating the need to protect and then de-protect the 4-hydroxy group:



Example II


N-(3-methoxy-4-hydroxybenzyl)-N'-octylthiourea was prepared by the following method.

N-3-methoxy-4-hydroxybenzyl)-N'-octylthiourea

10.6 ml of 5 N NaOH was added to a suspension of 5.0 g 3-methoxy-4-hydroxybenzylamine hydrochloride in 35 ml dimethylformamide (DMF). The resulting solution was cooled in an ice bath and 4.96 g of octyl isothiocyanate was added dropwise. The reaction mixture was stirred for six hours at room temperature, poured into 500 ml of H₂O, and then extracted with three successive 100 ml portions of diethyl ether. The combined ether extracts were returned to the separatory funnel and extracted with 75 ml 1 N HCl, 75 ml saturated NaHCO₃, 75 ml H₂O and 75 ml brine. The organic phase was dried over MgSO₄, filtered and evaporated. The crude product was crystallized from methanol yielding 4.58 gm of white solid M.P. 100—101°. Analysis calculated for C₁₇H₂₈N₂O₂S: C, 62.94; H, 8.70; N, 8.64; S, 9.88. Found: C, 63.05; H, 8.67; N, 8.66; S, 10.03.

Example III

N-(3-methoxy-4-hydroxybenzyl)-N'-dodecylthiourea was similarly prepared.



Analysis calculated for $C_{21}H_{35}N_2O_2S$: C, 66.20; H, 9.53; N, 7.33. Found: C, 66.33; H, 9.54; N, 7.33.

By use of the appropriate isocyanate or isothiocyanate in one of the foregoing syntheses, any and all of the compounds of this invention can be prepared.

5 C. Topical Compositions

The urea derivatives as previously described are useful when topically applied to skin. Compositions containing these compounds are also useful for topical application to skin. The compositions comprise a safe and effective amount, usually at least about 0.5%, preferably from about 1% to about 2%, of the urea derivative. The balance of the composition further comprises a
10 pharmaceutically acceptable carrier. Suitable carriers for the urea derivatives preferably remain in place on the skin as a continuous film and resist being washed off easily by perspiration or by immersion in water. Generally, the carrier is organic in feature and capable of having dispersed or dissolved therein the urea derivative. Lotions, creams, solutions, gels and solids are common physical forms of the compositions herein. A more detailed description of such forms follows.

15

1. Lotions

The lotions can comprise an effective amount of the urea derivative; from 1% to 25%, preferably from 3% to 15%, of an emollient; the balance being water, a C_2 or C_3 alcohol, or a mixture of water and the alcohol. Several emollients are known. Examples of such emollients are as follows:

20 1. Hydrocarbon oils and waxes. Examples are mineral oil, petrolatum, paraffin, ceresin, ozokerite, microcrystalline wax, polyethylene, and perhydrosqualene.

2. Silicone oils, such as dimethylpolysiloxanes, methylphenylpolysiloxanes, water-soluble and alcohol-soluble silicone-glycol copolymers.

3. Triglyceride fats and oils such as those derived from vegetable, animal and marine sources.
25 Examples include castor oil, safflower oil, cotton seed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, and soybean oil.

4. Acetoglyceride esters, such as acetylated monoglycerides.

5. Ethoxylated glycerides, such as ethoxylated glyceryl monostearate.

6. Alkyl esters of fatty acids having 10 to 20 carbon atoms. Methyl, isopropyl, and butyl esters of
30 fatty acids are useful herein. Examples include hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl, palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate.

7. Alkenyl esters of fatty acids having 10 to 20 carbon atoms. Examples thereof include oleyl
35 myristate, oleyl stearate, and oleyl oleate.

8. Fatty acids having 9 to 22 carbon atoms. Suitable examples include pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic, and erucic acids.

9. Fatty alcohols having 10 to 22 carbon atoms. Lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecyl alcohols are examples of
40 satisfactory fatty alcohols.

10. Fatty alcohol ethers. Ethoxylated fatty alcohols of 10 to 20 carbon atoms include the lauryl, cetyl, stearyl, isostearyl, oleyl, and cholesterol alcohols having attached thereto from 1 to 50 ethylene oxide groups or 1 to 50 propylene oxide groups, or a mixture thereof.

11. Ether-esters such as fatty acid esters of ethoxylated fatty alcohols.

45 12. Lanolin and derivatives. Lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated lanolin, ethoxylated lanolin alcohols, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin, acetylated lanolin alcohols, lanolin alcohols linoleate, lanolin alcohols ricinoleate, acetate of lanolin alcohols ricinoleate, acetate of ethoxylated alcohols-esters, hydrogenolysis of lanolin, ethoxylated hydrogenated lanolin, ethoxylated sorbitol lanolin, and
50 liquid and semisolid lanolin absorption bases are illustrative of emollients derived from lanolin.

13. Polyhydric alcohols and polyether derivatives. Propylene glycol, dipropylene glycol, polypropylene glycol (M.W. 2000—4000), polyoxyethylene polyoxypropylene glycols, polyoxypropylene polyoxyethylene glycols, glycerol, ethoxylated glycerol, propoxylated glycerol, sorbitol, ethoxylated sorbitol, hydroxypropyl sorbitol, polyethylene glycol (M.W. 2000—6000),
55 methoxy polyethylene glycols, 350, 550, 750, 2000, 5000, poly[ethylene oxide] homopolymers (M.W. 100,000—5,000,000), polyalkylene glycols and derivatives, hexylene glycol (2-methyl-2,4-pentanediol), 1,3-butylene glycol, 1,2,6-hexanetriol, ethohexadiol USP (2-ethyl-1,3-hexanediol), C_{15} — C_{18} vicinal glycol, and polyoxypropylene derivatives of trimethylolpropane are examples thereof.

14. Polyhydric alcohol esters. Ethylene glycol mono- and di-fatty acid esters, diethylene glycol
60 mono- and di-fatty acid esters, polyethylene glycol (M.W. 200—6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 monooleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty acid esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid ester, sorbitan fatty acid
65 esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters.

15. Wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate.
16. Beeswax derivatives, e.g. polyoxyethylene sorbitol beeswax. These are reaction products of beeswax with ethoxylated sorbitol of varying ethylene oxide content, forming a mixture of ether-esters.
17. Vegetable waxes including carnauba and candelilla waxes.
18. Phospholipids such as lecithin and derivatives.
19. Sterols. Cholesterol, cholesterol fatty acid esters are examples thereof.
20. Amides such as fatty acid amides, ethoxylated fatty acid amides, solid fatty acid alkanolamides.

The lotions further comprise from 1% to 10%, preferably from 2% to 5%, of an emulsifier. The emulsifiers can be nonionic, anionic or cationic. Examples of satisfactory nonionic emulsifiers include fatty alcohols having 10 to 20 carbon atoms, fatty alcohols having 10 to 20 carbon atoms condensed with 2 to 20 moles of ethylene oxide or propylene oxide, alkyl phenols with 6 to 12 carbon atoms in the alkyl chain condensed with 2 to 20 moles of ethylene oxide, mono- and di-fatty acid esters of ethylene glycol wherein the fatty acid moiety contains from 10 to 20 carbon atoms, fatty acid monoglycerides wherein the fatty acid moiety contains from 10 to 20 carbon atoms, diethylene glycol, polyethylene glycols of molecular weight 200 to 6000, propylene glycols of molecular weight 200 to 3000, glycerol, sorbitol, sorbitan, polyoxyethylene sorbitol, polyoxyethylene sorbitan, and hydrophilic wax esters. Suitable anionic emulsifiers include the fatty acid soaps, e.g. sodium, potassium and triethanolamine soaps, wherein the fatty acid moiety contains from 10 to 20 carbon atoms. Other suitable anionic emulsifiers include the alkali metal, ammonium or substituted ammonium alkyl sulfates, alkyl arylsulfonates, and alkyl ethoxy ether sulfonates having 10 to 30 carbon atoms in the alkyl moiety. The alkyl ethoxy ether sulfonates contain from 1 to 50 ethylene oxide units. Satisfactory cationic emulsifiers are the quaternary ammonium, morpholinium and pyridinium compounds. Certain of the emulsifiers described in preceding paragraphs also have emulsifying properties. When a lotion is formulated containing such an emollient, an additional emulsifier is not needed, though it can be included in the composition.

The balance of the lotion is water or a C₂ or C₃ alcohol, or a mixture of water and the alcohol. The lotions are formulated by simply admixing all of the components together. Preferably the urea derivative is dissolved in the mixture. Optional components such as common additives can be included. One common additive is a thickening agent at a level of from 1% to 10% of the composition. Examples of suitable thickening agents include: cross-linked carboxypolymethylene polymers, ethyl cellulose, polyethylene glycols, gum tragacanth, gum kharaya, xanthan gums and bentonite.

2. Creams

Compositions of this invention also can be formulated in a cream form. The creams comprise an effective amount of the urea derivative; from 5% to 50%, preferably from 10% to 25%, of an emollient; the balance being water. The emollients above described can also be used in the cream compositions. Optionally the cream form contains a suitable emulsifier, as previously described. When an emulsifier is included, it is in the composition at a level from 3% to 50%, preferably from 5% to 20%.

3. Solutions

The compositions of this invention can be also formulated in a solution form. The solution form comprises an effective amount of the urea derivative; the balance being a suitable organic solvent. Suitable organic materials useful as the solvent or a part of a solvent system are as follows: propylene glycol, polyethylene glycol (M.W. 200—600), polypropylene glycol (M.W. 425—2025), glycerine, sorbitol esters, 1,2,6-hexanetriol, ethanol, isopropanol, diethyl tartrate, butanediol, and mixtures thereof. Such solvent systems can also contain water.

These compositions in the solution form can be applied to the skin as is, or else can be formulated into an aerosol and applied to the skin as a spray-on. The aerosol compositions further comprise from 25% to 80%, preferably from 30% to 50%, of a suitable propellant. Examples of such propellants are the chlorinated, fluorinated and chlorofluorinated lower molecular weight hydrocarbons. Nitrous oxide, carbon dioxide, butane, and propane are also used as propellant gases. These propellants are used at a level sufficient to expel the contents of the container.

4. Gels

Compositions herein can be formulated into a gel form by simply admixing a suitable thickening agent to the previously described solution compositions. Examples of suitable thickening agents have been previously described with respect to the lotions.

The gelled compositions comprise an effective amount of the urea derivative; from 5% to 75%, preferably from 10% to 50%, of an organic solvent as previously described; from 0.5% to 20%, preferably from 1% to 10% of the thickening agent; the balance being water.

5. Solids

The compositions of this invention can also be formulated into a solid form. Such forms have use as a stick-type composition intended for application to the lips or other parts of the body. Such

compositions comprise an effective amount of the urea derivative and from 50% to 98%, preferably from 60% to 90%, of the previously described emollients. This composition can further comprise from 1% to 20%, preferably from 5% to 15%, of a suitable thickening agent, and optionally emulsifiers and water. Thickening agents previously described with respect to the gelled compositions are suitable

5 herein.

Additives commonly found in topical compositions such as preservatives, e.g., methyl and ethylparaben, dyes and perfume can be included in any of the previously described compositions.

D. Pharmaceutical Compositions and Dosage Forms for Systemic Administration

10 The urea derivatives of the present invention are also useful when used systemically, for example by parenteral administration. The dosage of the urea derivative which is both safe and effective to provide analgesic or anti-irritant activity will vary with the particular condition being treated, the severity of the condition, the duration of treatment, the specific urea derivative employed and its usage concentration, and like factors within the specific knowledge and expertise of the patient or the

15 attending physician and commensurate with a reasonable benefit/risk ratio associated with the use of any drug compound. The systemic dosages and dosage ranges given herein are based on delivery of the urea derivative to a 70 kg human and can be adjusted to provide equivalent dosages for patients of different body weights.

For mammals, especially humans, individual doses of from 0.1 mg to over 100 mg with total dosages of from 0.5 mg to 500 mg are acceptable. Individual doses of from 0.5 mg to 50 mg with total dosages of from 1 mg to 100 mg are preferred. Individual doses of from 5 mg to 25 mg with total dosages of from 10 mg to 50 mg are especially preferred. While dosages higher than the foregoing are effective, toxicity and side effects will present problems in some individuals.

The urea derivatives can be administered parenterally in combination with a pharmaceutically acceptable carrier such as corn oil, Cremophor EL, or sterile, pyrogen-free water and a water-miscible solvent (e.g. ethyl alcohol) at a practical amount of the urea derivative per dose. Parenteral administration can be by subcutaneous, intradermal, intramuscular, intraarticular, or intravenous injection. The dosage by these modes of administration is usually in the range of from about 10 mg to about 500 mg per day.

30 As used herein the term "pharmaceutically acceptable carrier" denotes a solid or liquid filler, diluent, or encapsulating substance. Some examples of substances which can serve as pharmaceutical carriers for the urea derivatives include: sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives, such as sodium carboxymethylcellulose, ethyl cellulose, cellulose acetate, powdered tragacanth; malt; gelatin; talc, stearic acid, magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; sugar; alginic acid; pyrogen-free water; isotonic saline; phosphate buffer solutions; cocoa butter (suppository base), emulsifiers, such as the Tweens^a as well as other non-toxic compatible substances typically used on pharmaceutical formulations. Wetting agents and lubricants such as sodium lauryl

40 sulfate, as well as colouring agents, and preservatives, can also be present. The pharmaceutical carrier employed in conjunction with the urea derivative is used at a concentration sufficient to provide a practical size to dosage relationship. Preferably, the pharmaceutical carrier comprises at least about 98% by weight of the total composition.

45 E. Methods for Providing Analgesia and Anti-irritancy

The present invention encompasses methods for providing analgesia or anti-irritancy in humans or lower animals in need thereof by administering a safe and effective amount, usually from about 10 mg to about 500 mg per patient per day, of the urea derivative, usually as a composition with a pharmaceutically acceptable carrier. The urea derivatives and compositions containing same can be administered by topical application or systemic administration. The urea derivatives and compositions containing same can be used to treat pain and to provide an analgesic effect in various disorders at the deeper structures, muscles, tendons, bursa and joints associated with disease and trauma, and in various other conditions in which compounds such as aspirin and morphine have heretofore been used to alleviate pain and discomfort.

55 A "stairstep" dose pattern of sequentially increasing doses is useful for intraperitoneal administration in overcoming temporary side effects and in administering larger doses of the urea derivatives of this invention. Thus, the initial dose should be selected to provide minimal side effects, and subsequent doses should be increased until the desired level of activity is obtained while still minimizing side effects. In general, an initial dose of 2—10 mg/kg can be conveniently used, followed by doubling of each subsequent dose until the desired level is reached. The doses are preferably separated by a period of at least two hours.

F. Depilatory Compositions Having Reduced Dermal Irritation and Methods for Reducing or Preventing Irritation Caused by Depilatory Agents

65 The present invention also encompasses depilatory compositions having reduced dermal

irritation, which comprise: a) a depilatory amount (e.g. at least about 2%) of a thioglycolate depilatory agent; and b) the urea derivative in an amount effective to reduce the dermal irritation caused by the thioglycolate depilatory agent. The depilatory compositions can be formulated using the pharmaceutically acceptable carriers previously described for topical compositions.

5 The present invention further encompasses a method for preventing or reducing the dermal irritation caused by a thioglycolate depilatory agent, which comprises the step of applying to at least a portion of a depilated area the urea derivative (or composition containing same) in an amount effective to prevent or reduce the irritation caused by treatment of the depilated area with the thioglycolate depilatory agent.

10 By "thioglycolate depilatory agent" is meant thioglycolic acid, its alkali metal, alkaline earth metal, and ammonium salt(s), or mixtures of the acid and its salt(s).

By "an amount effective to prevent or reduce irritation" is meant an amount of the urea derivative (or composition containing same) effective to reduce or prevent the irritation caused by treatment of the depilated area with the thioglycolate depilatory agent at a reasonable benefit/risk ratio. The amount of the urea derivative used will vary with the severity of the irritation, the duration of the treatment, the specific formulation employed, the concentration of the urea derivative therein, and like factors.

15 By "depilated area" is meant that area which is, or is about to be, depilated by treatment with a thioglycolate depilatory agent.

By the term "applying" with regard to preventing or reducing depilatory irritation is meant the direct laying on or spreading of the urea derivative (including compositions containing same) on skin tissue which is, or is about to be depilated. The urea derivative can be applied before and/or after treatment of the depilated area with the thioglycolate depilatory agent to prevent or reduce irritation caused thereby. Application of the urea derivative to the depilated area after treatment with the depilatory agent is preferred, especially when lower concentrations of the urea derivative are used. The number of applications needed to provide effective irritation prevention or reduction can depend upon the concentration of the urea derivative used, and when the urea derivative is applied in relation to the treatment with the depilatory agent. Application of the urea derivative soon after depilation, e.g. within about 6 to about 12 hours, provides effective irritation prevention or reduction, especially in conjunction with additional applications on subsequent days. Multiple applications (2 or more sequential, time spaced applications) soon after depilation are particularly effective. The length of time during which the urea derivative is left on the depilated area can also determine its effectiveness. An application duration of at least about 1 hour, preferably from about 1 to about 2 hours, should provide effective irritation prevention or reduction.

Specific Embodiments of Compositions Containing N-(3-Methoxy-4-Hydroxybenzyl and phenyl) Ureas and Thioureas

35 The following examples illustrate compositions within the scope of the present invention, but are not limiting thereof.

Embodiment I

Lotion

40	Isopropyl myristate	8%
45	Corn oil	5%
	Propylene glycol	5%
50	Triethanolamine oleate	5%
	Xanthan gum	0.5%
	N-(3-methoxy-4-hydroxybenzyl)-N'-hexylurea	0.5%
55	Water	Balance

0 068 590**Embodiment II****Cream**

5	Isopropyl myristate	5%
	Sorbitol	
	Propylene glycol	10%
10	Triethanolamine stearate	17%
	N-(3-methoxy-4-hydroxybenzyl)-N'-dodecylthiourea	1%
15	Water	Balance

Embodiment III**Gel**

20	Oleyl alcohol	1%
	Propylene glycol	20%
25	Triethanolamine	0.5%
	Ethanol	57%
30	Carbomer 940	0.5%
	N-(3-methoxy-4-hydroxybenzyl)-N'- <i>cis</i> -12-docosenylurea	0.5%
35	Water	Balance

Embodiment IV**Solution**

40	Propylene glycol	10%
	Polyethylene glycol 400	2%
45	Tween 80 ^R	1%
	Ethanol	50%
50	N-(3-methoxy-4-hydroxyphenyl)-N'-hexenylthiourea	1%
	Water	Balance

Embodiment V**Ointment**

55	Oleyl alcohol	alcohol	30%
60	Cetyl alcohol		40%
	Propylene glycol		28%
65	N-(3-methoxy-4-hydroxyphenyl)-N'- <i>cis</i> -12-hexadecenylurea		2%

Effectiveness of N-(3-Methoxy-4-Hydroxybenzyl and phenyl) Ureas and Thioureas in Producing Analgesia and in Reducing or Preventing Depilatory Irritation

A. Analgesia

1. Mouse Hot Plate Tests

The Mouse Hot Plate (MHP) model system is designed to detect and evaluate agents which elevate the threshold for the perception of pain. Classically, this method has been utilized primarily to evaluate narcotic type analgesic agents such as morphine. Unless administered in toxic quantities, non-steroidal anti-inflammatory agents such as aspirin or acetaminophen exhibit little or no activity in this system.

Male CF-1 mice were used. Animals were divided into groups of 8 to 12 and then tested on the "hot plate" to determine their predrug response time which was 4.9 seconds (average). Each animal was then treated with either a test composition (0.5% to 2% of the particular urea or thiourea in an isotonic saline solution containing ethyl alcohol and Tween® 80) or a control composition (same as test composition but without the urea or thiourea). Treatment was by injection prior to test initiation.

2. Procedure

The mice were placed on a 55°C heated surface and their responses were observed. The endpoint is either a rapid fanning or licking of a paw. To prevent physical injury, the animals were not allowed to remain in contact with the heated surface for more than 60 seconds. The exposure time required to elicit the endpoint response is a measure of the pain threshold. The difference in time required to elicit the endpoint response before and after treatment provides a measure of analgesia. The increase in time required to elicit the endpoint response in treated animals versus control composition is a second measure.

3. Results

The results from the MHP testing of the N-(3-methoxy-4-hydroxybenzyl or phenyl) ureas or thioureas were as follows:

TABLE I

Compound	Method**	Dosage*** (mg/kg)	Post-Drug Response (sec)
Aspirin*	O	360	6.0
Acetaminophen*	O	450	5.3
Morphine Sulfate*	IP	13	13.1
Morphine Sulfate*	IP	25	17.4
Control	SC	—	5.0
<u>N-(3-methoxyl-4-hydroxybenzyl)</u>			
Octylurea	SC	0.68 mM/kg	6.5
Octylthiourea	SC	0.68 mM/kg	14.7
Dodecylthiourea	SC	0.68 mM/kg	6.5
Phenylthiourea	SC	0.68 mM/kg	8.0
4-fluorophenylthiourea	SC	0.68 mM/kg	6.4
1,1,3,3-tetramethylbutylthiourea	IP	25	7.0

0068 590

TABLE I (continued)

Compound	Method**	Dosage*** (mg/kg)	Post-Drug Response (sec)
<u>N-(3-methoxy-4-hydroxyphenyl)</u>			
5 Octylurea	SC	4,8,15,25,50	6.2
Octylurea	IP	4,8,15,25	11.5
10 Octylurea	SC	50	9.8
	IP	50	52.8****
Octylthiourea	SC	0.68 mM/kg	9.5
15			

* Commercial Analgesics

** O — oral; SC — subcutaneous; IP — intraperitoneal

*** Multiple doses were 2 hours apart

20 **** IP dose was 24 hrs. after SC dose; 40% of animals dead

The data in Table I shows that both N - (3 - methoxy - 4 - hydroxy - benzyl) - N' - octylthiourea and -phenylthiourea, as well as N - (3 - methoxy - 4 - hydroxyphenyl) - N' - octylurea and octylthiourea, have significant analgesic activity relative to the control composition and the commercial analgesics.

25

B. Preventing or Reducing Depilatory Irritation

Groups of 8 male Sprague-Dawley rats weighing 90—115 grams were used for testing the effectiveness of the N-(3-methoxy-4-hydroxybenzyl or phenyl) urea and thioureas of the present invention in preventing or reducing depilatory irritation. The animals were clipped and depilated with 30 Nair®, a commercially available thioglycolate depilatory. The test compositions (2% of the urea or thiourea in an isotonic saline solution containing 48% ethyl alcohol, 4% Tween® 80) or a control composition (same as test composition but without the urea or thiourea) were applied to one quadrant (Treated Area) of the depilated area once, 2 hours after depilation on first day; four times, 2 hours apart on second day; and three times, 2 hours apart on third day for a total of eight applications. The duration 35 of each application was 2 hours. The remaining three quadrants of the depilated area was left untreated (Untreated Area). Oral ingestion was prevented by the use of "Elizabethan" collars. On the fourth day, the animals were depilated a second time and evaluated for irritation four hours later.

Irritation scores for each animal were determined by visual inspection using the following subjective evaluation scale:

40	Score	Description of Irritation
	0	No irritation
45	0.5	No scab formation, faint white scale
	1.0	Scab formation (pale orange/orange) over less than 10% of area
	2.0	Mild to moderate intensity scab formation (pale orange) over 10—33% of area
50	3.0	Mild to moderate intensity scab formation (pale orange) over 33—75% of area
	3.5	Moderate intensity scab formation (pale orange, occasional deep orange/red) over 75—90% of area
55	4.0	Moderate to severe intensity scab formation (deep orange, occasional pale orange) over 90—100% of area
	5.0	Severe intensity scab formation (deep orange/red) over 100% of area

60

The irritation scores were totaled for each of the 8-animal groups, the maximum cumulative score being 40. A cumulative score of less than 8 indicated a minimal level of irritation; a score of 8—24 indicated a higher, but acceptable level of irritation; a score of above 24 indicated an unacceptable level of irritation. The results from this testing of the N-(3-methoxy-4-hydroxybenzyl or phenyl) ureas and

65 thioureas were as follows:

TABLE II


5	Composition	Cumulative Score	
		Treated Area	Untreated Area
	Control	26.5	20
10	N-(3-methoxy-4-hydroxyphenyl)- octyl thiourea	15	6
	octyl urea	16	9
15	N-(3-methoxy-4-hydroxybenzyl)*		
	1,1,3,3-tetramethylbutylthiourea	1.3	13.3
20	phenylthiourea	15.3	30.0

*extrapolated from group of 6 animals

The data in Table II shows that N - (3 - methoxy - 4 - hydroxyphenyl) - N' - octylurea and octylthiourea, and N - (3 - methoxy - 4 - hydroxybenzyl) - N' - 1,1,3,3 - tetramethylbutylthiourea, were effective in preventing or reducing depilatory irritation in both the Treated and Untreated Areas; the N - (3 - methoxy - 4 - hydroxybenzyl) - N' - phenylthiourea was effective in the Treated Area.

Claims

1. A compound characterized in that it has the formula:



wherein X is O or S, wherein $n = 1$ or 0 , wherein R is $\text{C}_3\text{--C}_{12}$ alkyl or aralkyl, branched or unbranched, aryl or haloaryl, or $\text{C}_3\text{--C}_{22}$ alkenyl, branched or unbranched, wherein R_1 is H, OH or OCH_3 , and R_2 is H or OH when $n = 1$, at least one of R_1 and R_2 being OH or OCH_3 , and wherein R_1 is OH or OCH_3 and R_2 is OH when $n = 0$; and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1 characterized in that it is the sodium, potassium, calcium, magnesium, or ammonium salt.

3. The compound or salt of Claim 1 or 2 characterized in that R_1 is OCH_3 and R_2 is OH.

4. The compound or salt of any of Claims 1—3 characterized in that X is S.

5. The compound or salt of any of Claims 1—4 characterized in that $n = 1$.

6. The compound or salt of any of Claims 1—5 characterized in that R is $\text{C}_6\text{--C}_{12}$ alkyl or aralkyl, or $\text{C}_6\text{--C}_{22}$ alkenyl.

7. The compound or salt of any of Claims 1—6 characterized in that R is 1,1,3,3-tetramethylbutyl or n-octyl.

8. A pharmaceutical composition in unit dosage form having analgesic activity characterized in that it comprises a safe and effective amount of the compound or salt of Claim 1, 2, 3, 4, 5, 6 or 7; and a pharmaceutically acceptable carrier.

9. A depilatory composition having reduced dermal irritation characterized in that it comprises:


a) a depilatory amount of a thioglycolate depilatory agent, and

b) the compound or salt of any of Claims 1—7 in an amount effective to prevent or reduce the dermal irritation of the thioglycolate depilatory agent.

10. The depilatory composition of Claim 9 characterized in that it further comprises a pharmaceutically acceptable carrier.

Patentansprüche


1. Eine Verbindung, dadurch gekennzeichnet, daß sie die Formel



- 15 aufweist, worin X O oder S ist, worin $n = 1$ oder 0 ist, worin R eine verzweigte oder unverzweigte C_3-C_{12} -Alkylgruppe oder Aralkylgruppe, Arylgruppe oder Halogenarylgruppe oder eine verzweigte oder unverzweigte C_3-C_{22} -Alkenylgruppe ist, worin R_1 H, OH oder OCH_3 ist und R_2 H oder OH ist, wenn $n = 1$ ist, wobei mindestens einer der Reste R_1 und R_2 OH oder OCH_3 bedeuten, und worin R_1 OH oder OCH_3 ist und R_2 OH ist, wenn $n = 0$ ist; und pharmazeutisch verträgliche Salze davon.
- 20 2. Die Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß sie das Natrium-, Kalium-, Calcium-, Magnesium- oder Ammoniumsalz ist.
3. Die Verbindung oder das Salz nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß R_1 OCH_3 ist und R_2 OH ist.
4. Die Verbindung oder das Salz nach einem der Ansprüche 1 bis 3, dadurch gekennzeichnet, daß
- 25 X S ist.
5. Die Verbindung oder das Salz nach einem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß $n = 1$ ist.
6. Die Verbindung oder das Salz nach einem der Ansprüche 1 bis 5, dadurch gekennzeichnet, daß R eine C_8-C_{12} -Alkyl- oder Aralkylgruppe oder eine C_8-C_{22} -Alkenylgruppe ist.
- 30 7. Die Verbindung oder das Salz nach einem der Ansprüche 1 bis 6, dadurch gekennzeichnet, daß R die 1,1,3,3-Tetramethylbutylgruppe oder die n-Octylgruppe ist.
8. Pharmazeutische Zubereitung in Einheitsdosierungsform mit schmerzlindernder Wirksamkeit, dadurch gekennzeichnet, daß sie eine sichere und wirksame Menge der Verbindung oder des Salzes von Anspruch 1, 2, 3, 4, 5, 6 oder 7 und einen pharmazeutisch verträglichen Träger enthält.
- 35 9. Enthaarungsmittel mit verringerter Hautreizung, dadurch gekennzeichnet, daß es enthält:
- a) eine enthaarende Menge eines Thioglycolat-Enthaarungsmittels und
- b) die Verbindung oder das Salz nach einem der Ansprüche 1 bis 7 in einer Menge, die wirksam ist, um die Hautreizung des Thioglycolat-Enthaarungsmittels zu verhindern oder zu verringern.
10. Enthaarungsmittel nach Anspruch 9, dadurch gekennzeichnet, daß es weiterhin einen pharmazeutisch verträglichen Träger enthält.
- 40

Revendications

1. Composé caractérisé en ce qu'il a la formule:



- où X est O ou S, où $n = 1$ ou 0, où R est un radical alkyle en C_3 à C_{12} ou aralkyle, ramifié ou non ramifié, aryle ou halogénaryle, ou un radical alcényle en C_3 à C_{22} , ramifié ou non ramifié, où R_1 est H, OH ou OCH_3 et R_2 est H ou OH quand $n = 1$, au moins l'un des radicaux R_1 et R_2 étant OH ou OCH_3 , et où R_1 est OH ou OCH_3 et R_2 est OH quand $n = 0$; et ses sels pharmaceutiquement acceptables.
- 60 2. Composé selon la revendication 1, caractérisé en ce qu'il est le sel de sodium, de potassium, de calcium, de magnésium ou d'ammonium.
3. Composé ou sel selon la revendication 1 ou 2, caractérisé en ce que R_1 est OCH_3 et R_2 est OH.
4. Composé ou sel selon l'une quelconque des revendications 1 à 3, caractérisé en ce que X est S.
- 65 5. Composé ou sel selon l'une quelconque des revendications 1 à 4, caractérisé en ce que $n = 1$.

6. Composé ou sel selon l'une quelconque des revendications 1 à 5, caractérisé en ce que R est un radical alkyle en C₆ à C₁₂ aralkyle ou alcényle en C₃ à C₂₂.

7. Composé ou sel selon l'une quelconque des revendications 1 à 6, caractérisé en ce que R est le radical 1,1,3,3-tétraméthylbutyl ou n-octyl.

6 8. Composition pharmaceutique sous forme de dose unitaire, ayant une activité analgésique, caractérisée en ce qu'elle comprend une quantité non toxique efficace du composé ou sel selon l'une des revendications 1, 2, 3, 4, 5, 6 ou 7; et un excipient pharmaceutiquement acceptable

9. Composition dépilatoire entraînant une irritation réduite du derme, caractérisée en ce qu'elle comprend a) une quantité dépilatoire d'un agent dépilatoire de type thioglycolate et b) le composé ou
10 sel selon l'une quelconque des revendications 1 à 7, en une quantité efficace pour prévenir ou réduire l'irritation du derme due à l'agent dépilatoire de type thioglycolate.

10. Composition dépilatoire selon la revendication 9, caractérisée en ce qu'elle comprend en outre un excipient pharmaceutiquement acceptable.

15

20

25

30

35

40

45

50

55

60

65